


Development of Pentose Fermenting Yeast Strains for Ethanol Production from Lignocellulosic Biomass

Maurizio Bettiga
Department of Applied Microbiology, Lund University
Lund, Sweden




LUND
UNIVERSITY


Fermentation of Lignocellulosic Hydrolysate



Saccharomyces cerevisiae Is a Good Choice



New Pentose Fermenting Strains



Available pathways

BACTERIAL
FUNGAL


New strain with the required pathways

Pentose fermenting strain?




A Combined Strategy to Reach the Goal

Metabolic systems work in a highly coordinated status




The Fitness Landscape



- Wright (1982) Evolution 36:427.
- Kauffman (1993) The origins of order. Oxford University press.



Variation Scans the Fitness Landscape



**Evolutionary engineering/mutagenesis
(Scan the fitness landscape)**

**Increased mutational rate:
Increased driving force through
the fitness landscape**

**Genetic engineering
(Jump)**


**Metabolic system in
"suboptimal" state**

Point 0: No pentose genes

Xylose Fermenting Strain



UNIVERSITEIT·STELLENBOSCH·UNIVERSITY
jou kennisvennoot·your knowledge partner



- Wahlbom et al (2003) FEMS Yeast Res 3:319-326
- Wahlbom et al (2003) Appl Environ Microbiol 69:740-746

(Proteome under study)
Karhumaa, submitted




LUND
UNIVERSITY


Performances in Operational Conditions

SSF, Fed Batch, Corn Stover

Xylose is co-consumed with glucose




By product (Xylitol) production is low



Lower Xylitol Production May Result from:

Increased Overall Metabolic Flux by Presence of Glucose




• Karhumaa et al (2006)
Appl Microbiol Biotechnol. In press.

- Jeffries et al (1984) Biotechnol Bioeng 27:171
- Jin et al (2004) Appl Env Microbiol 70:6816-25



Lower Xylitol Production May Result from:

Presence of electron acceptors
(Pretreatment by-products)




Arabinose Fermenting Strain/1




Genetic
engineering

AraA/B/D

AraA; AraD: multiple copy integration
AraB: single copy integration




Proof of Principle




- Nilsson et al (2005) Appl Env Microb. 71:7866-71
- Petersson et al (2006) Yeast 23:455-64

Reduction Of Furans Improves Tolerance

HMF
5-hydroxymethyl furfural




HMF-OH




NAD(P)H


furfural




F-OH



NAD(P) +



Tolerance⁺ strain



- Nilsson et al (2005) Appl Env Microb. 71:7866-71
- Wahlbom and Hahn-Hägerdal (2002) Biotechnol Bioeng 78:172-78.

Identification of HMF-reducing Enzymes/1

NAD(P)H-dependent activity

1


Microarray analysis

Non-Tolerant Vs Tolerant strain
Genes Induced?


2

Over expression of best candidates

HMF-reducing activity (*in vitro*)



HMF Uptake (*in vivo*)



ADH6

Acknowledgments

Marie Gorwa-Grauslund

Bärbel Hahn-Hägerdal

Applied Microbiology




Kaisa Karhumaa

Violeta Sanchez Nogue

Rosa Garcia Sanchez

João Ricardo Moreira de Almeida



<http://www.tmb.lth.se>



LUND
UNIVERSITY